

# Hip and Knee Surgical Care Path

## Release Date: 2024

Quick  
links:


- |  |   |
|--|---|
| <a href="#">Abbreviations</a>  | <a href="#">Care Path Diagram</a>                         |
| <a href="#">Referral to Assessment to Surgical/Medical/Functional Optimization</a> | <a href="#">Same Day Admit to Intra-Operative Surgery</a> |
| <a href="#">Acute Post-Operative Care</a>  | <a href="#">Recovery</a>                                  |
| <a href="#">Subacute</a>   | <a href="#">References</a>                                |

This care path guides the collaborative workflow between the interdisciplinary teams at hospitals and central intake clinics in Alberta providing hip and knee arthroplasty services. It is provincially endorsed encompassing instructions and performance indicators that are evidence-based and derived from clinical consensus. The care path has demonstrated that an evidence-based, collaborative, and integrated approach to surgical management has positive health system and patient outcomes.

## Abbreviations

Term	Meaning	Term	Meaning
ADL	Activities of Daily Living	NSAID	Non-steroid anti-inflammatory
AHS	Alberta Health Services	OR	Operative Room
AP	Anterior-posterior	OSA	Obstructive Sleep Apnea
ASA	American Society of Anesthesiologists	OT	Occupational Therapist
Bid	<i>Bis in die</i> – twice daily	PCN	Primary Care Network
BMI	Body Mass Index	PCP	Primary Care Physician
BPMH	Best Possible Medication History	prn	<i>Pro re nata</i> – take as needed
CBC	Complete Blood Count	PO	<i>Per os</i> – by mouth
CHG	Chlorhexidine Gluconate	POD	Post operative days
CM	Case Manager	PONV	Post operative nausea and vomiting
DAT	Diet as Tolerated	PT	Physiotherapist
DB & C	Deep breath and cough	Pt	Patient
D/C	Discharge	q_h	<i>Quaque hora</i> - every _ hours
DOAC	Direct Oral Anticoagulants	RN	Registered Nurse
DOS	Duration of stay: 24 hours = <sup>1</sup> day; defined as from when patient checks in for surgery to when (s)he leaves the hospital	ROM	Range of Motion
DVT	Deep Vein Thrombosis	RR	Respiratory Rate
ECG	Electrocardiogram	SBP	Systolic Blood Pressure
EQ5D	euroQol – 5 dimension – 5 levels	THA	Total Hip Arthroplasty
F/A	Foot and Ankle	TKA	Total Knee Arthroplasty
GI	Gastro-intestinal	TXA	Tranexamic Acid
GP	General Practitioner	QID	<i>Quater in Die</i> – four times per day
Hgb	Hemoglobin	WBC	White Blood Cells
IM	Internal Medicine	Wt	Weight
INR	International Normalized Ratio	WOMAC	Western Ontario and McMaster Universities Osteoarthritis Index
IPA	Isopropyl alcohol		
IV	Intravenous		
LMWH	Low Molecular Weight Heparin		
MSK	Musculoskeletal Specialist		

# The Hip and Knee Arthroplasty Care Path




### Legend

- Clinic – Activity
- Clinic – Decision point
- Clinic – Decision point Non-Surgical – refer to Conservative Care Path when available
- Hospital – Same Day Admit to Intra - Operative Surgery
- Hospital – Acute Post-Operative Care

## SECTION

# Referral to Assessment to Surgical/Medical/Functional Optimization



**Pre-referral and Referral**

Guidelines	Suggested Tools
<ul style="list-style-type: none"> <li>❑ <b>For Authorized Practitioner:</b> <ul style="list-style-type: none"> <li>• All referrals sent to Zone FAST Office</li> <li>• FAST forwards referrals that are complete and appropriate to designated Hip and Knee Replacement Clinic</li> <li>• Authorized practitioner submits completed referral form including diagnostic imaging and applicable consult reports</li> <li>• Referring authorized practitioner designates preferred Surgeon or next available Surgeon</li> <li>• Referring authorized practitioner will be contacted if preferred Surgeon is not available within appropriate wait standards.</li> </ul> </li> <li>❑ <b>For Primary Assessments:</b> <ul style="list-style-type: none"> <li>• All referrals triaged in a Hip and Knee Replacement Clinic</li> <li>• All referrals receipt acknowledged within 7 working days, and screened for appropriateness within 14 days<sup>1</sup></li> <li>• Triaging:                             <ol style="list-style-type: none"> <li>1. Referral is <u>inappropriate or incomplete</u>: send back to FAST to notify practitioner</li> <li>2. Referral is <u>appropriate and accepted</u>: notice of acceptance sent to practitioner by clinic                                     <ol style="list-style-type: none"> <li>a. Pts that require an MSK assessment to determine optimal treatment, are booked for an MSK Assessment Visit (see below)</li> <li>b. Otherwise Pt is booked for Initial Surgical Consult (see below)</li> </ol> </li> </ol> <p>Note: Appropriate and accepted referrals <i>should</i> be booked to receive a clinic evaluation within 40 working days<sup>2</sup></p> </li> <li>• If accepted, Pt instructions communicated to each Pt                             <ul style="list-style-type: none"> <li>▪ Pt advised to select “Buddy”/Family member to attend clinic visits</li> </ul> </li> </ul> </li> <li>❑ <b>For Revision:</b> <ul style="list-style-type: none"> <li>• Monitored Pt booked by Surgeon who did primary; or</li> <li>• Monitored Pt referred to revision specialist by Surgeon who did primary; or</li> <li>• Non-monitored Pt’s authorized practitioner completes referral</li> <li>• Pt advised to select “Buddy”/Family member to attend clinic visits</li> </ul> </li> </ul>	<p style="background-color: #0056b3; color: white; padding: 2px;"><b>Required for Referral Tests/Diagnostics</b></p> <ul style="list-style-type: none"> <li>❑ <b>Provincial Adult Orthopedic &amp; Spine Referral Pathways</b></li> <li>❑ <b>Knee:</b> <ul style="list-style-type: none"> <li>• AP WB</li> <li>• Lateral</li> <li>• Skyline</li> <li>• Rosenberg</li> </ul> </li> <li>❑ <b>Hip:</b> <ul style="list-style-type: none"> <li>• AP pelvis centered at pubis</li> <li>• AP and lateral of proximal half of affected femur</li> </ul> </li> <li>❑ <b>For Revision:</b> <ul style="list-style-type: none"> <li>• Use monitored images if completed</li> <li>• If monitoring images have not been done, then complete full set as above shoot through lateral of hip if requested</li> </ul> </li> </ul>

## Hip and Knee Clinic Evaluation and Detailed Assessment

MSK Assessment Visit	Suggested Tools
<ul style="list-style-type: none"> <li><input type="checkbox"/> <b>Preparation:</b> <ul style="list-style-type: none"> <li>• Pt completes self assessment</li> <li>• Outcome measurement tool completed on all Pts</li> </ul> </li> <li><input type="checkbox"/> <b>During MSK Assessment:</b> <ul style="list-style-type: none"> <li>• Pt validate referral package material</li> <li>• MSK specialist/Surgeon decides whether a surgical assessment is required or the Pt is non-surgical, see criteria (right)                             <ol style="list-style-type: none"> <li>1. Pt has maximized conservative treatment and a surgical assessment is required</li> <li>2. Pt has not maximized conservative treatment – conservative treatments can still be exploited to improve quality of life and function (Pt. is non-surgical, see below)</li> <li>3. Pt is not an appropriate candidate for an elective surgery (see criteria, right). Conservative treatments can be explored (Pt. is non-surgical, see below)</li> </ol> </li> <li>• If applicable, subsequent appointments are booked (see below)</li> </ul> </li> <li><input type="checkbox"/> <b>If 2<sup>nd</sup> Opinion is Required:</b> <ul style="list-style-type: none"> <li>• 2<sup>nd</sup> opinion can be requested by referring authorized practitioner or Pt.</li> <li>• Referring authorized practitioner to submit new Referral and indicate 2<sup>nd</sup> opinion</li> <li>• Pts to call clinic directly and request 2<sup>nd</sup> opinion</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li><a href="#">Hip and Knee Outcomes Tool (Combined WOMAC and EQ5D-5L)</a></li> <li><a href="#">OA Self Management Toolkit</a></li> </ul> <hr/> <p><b>Criteria for Non-Surgical Pt</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> Cognitive/neurologic impairment (Surgeon discretion)</li> <li><input type="checkbox"/> Orthopaedic challenges such as history of infection, or technical infeasible, or the joint cannot be reconstructed</li> <li><input type="checkbox"/> Pt refuses surgery</li> <li><input type="checkbox"/> Not willing to be compliant with the care path</li> <li><input type="checkbox"/> Extreme medical risk</li> </ul>

### Non-Surgical Patient

Non-Surgical Care
<ul style="list-style-type: none"> <li><input type="checkbox"/> <b>If Pt is Designated as Non-Surgical:</b> <ul style="list-style-type: none"> <li>• MSK specialist/RN or Surgeon/RN/CM completes either Surgical Optimization Pt Plan or Non-Surgical Pt Plan</li> <li>• Some non-surgical Pts return to clinic for non-surgical treatment. Non-surgical Pts who return for treatment book with clinic representative</li> <li>• Decision and accompanying documentation communicated back to referring authorized practitioner</li> <li>• Follow 2<sup>nd</sup> Opinion steps (above) if required</li> </ul> </li> </ul>
Suggested Tools
<p>➤ <a href="#">Please also refer to OA Conservative Management Care Map – Developed by the Conservative OA Clinical Committee</a></p>

**Surgical Assessment**

Initial Surgical Consult	Additional Tests and Diagnostic Imaging
<p><input type="checkbox"/> <b>Preparation:</b></p> <ul style="list-style-type: none"> <li>• All surgical Pt completes Hip/Knee Patient Reported Outcome and Experience Measures</li> <li>• All surgical Pts assigned to a Hip and Knee Replacement Surgeon and CM based on next available or requested Surgeon (if within standard waits)</li> </ul> <p><input type="checkbox"/> <b>During Hip and Knee Replacement Clinic Team Evaluation:</b></p> <ul style="list-style-type: none"> <li>• Surgeon and CM validate/complete referral package material (history and demographics)</li> <li>• X-rays ordered for views missing from referral stage</li> <li>• Surgeon assess risk factors and determines feasibility of surgery:               <ol style="list-style-type: none"> <li>1. Pt has too many risk factors and Surgeon is unwilling to proceed (Pt is non-surgical, see above)</li> <li>2. Pt still has conservative measures they can attempt (Pt is non-surgical, see above)</li> <li>3. Pt is suitable for surgery, but existing risk factors will require optimization and management (follow Surgical Pt Optimization, below)</li> <li>4. Pt is suitable for surgery and is ready for teaching (follow Surgical Pt Optimization, below)</li> <li>5. Consider if Pt is day surgery candidate (refer to Exclusion Criteria for Day Surgery Candidacy)</li> </ol> </li> <li>• Start surgical Pt agreement<sup>3</sup></li> <li>• Order minor-treatments, pre-op PT, clinic OT consult, nutrition consult, home visit and referrals for medical clearance and peri-operative blood consultation (see Surgical Pt Optimization, below, for criteria)</li> <li>• Surgical Pts advised dental clearance is required, at Surgeon discretion</li> <li>• CM reviews consult referrals and lab/imaging requisitions with Pt and advises Pt of timelines               <ol style="list-style-type: none"> <li>1. Copy of plan given to Pt</li> <li>2. Copy of plan sent to referring authorized practitioner and PCP, if different, within 5 working days of Pt's consult</li> </ol> </li> </ul>	<p><input type="checkbox"/> <b>For Knees</b></p> <ul style="list-style-type: none"> <li>• Three foot standing of limb at Surgeon's discretion</li> </ul> <p><input type="checkbox"/> <b>For Hips</b></p> <ul style="list-style-type: none"> <li>• Additional film as required at Surgeon's discretion</li> </ul> <p>Note: Pre-operative x-rays with templating spheres done at the discretion of the Surgeon</p>
	<b>Teaching/Discharge</b>
	<p><input type="checkbox"/> Surgical Pts are given Pt education package (guidebook and resources) to review at home</p> <p><input type="checkbox"/> At clinic's discretion, surgical Pts are directed:</p> <ul style="list-style-type: none"> <li>• to local pharmacy for full medication review, and</li> <li>• for initiation of tobacco cessation program, and/or alcohol cessation program, if applicable.</li> </ul> <p>Note: Pt to bring all documentation from complementary programs to next clinic appointment.</p> <p><input type="checkbox"/> Surgical Pts are encouraged to visit their PCP to develop their goals of care and personal directives</p>
	<b>Suggested Tools</b>
	<p><a href="#">Hip and Knee Outcomes Tool (Combined WOMAC and EQ5D-5L)</a></p> <p><input type="checkbox"/> <a href="#">Surgical Patient Agreement<sup>3</sup></a></p> <p><input type="checkbox"/> <a href="#">Buddy Agreement</a></p> <p><input type="checkbox"/> <a href="#">Canadian Nutrition Screening Tool</a></p>
	<b>Patient/Family/Buddy Responsibility</b>
	<p><input type="checkbox"/> Select "Buddy" to attend all clinic visits, especially teaching session</p> <p><input type="checkbox"/> Buddy to sign agreement of duties throughout surgery and discharge, and a Plan B if not available on surgery dates</p> <p><input type="checkbox"/> Buddy to notify Hip and Knee Replacement CM if Pt's medical status changes</p>

## Surgical Patient Optimization

### Guidelines

Optimization Management	ASA Classification														
<ul style="list-style-type: none"> <li><input type="checkbox"/> Pt will not proceed to surgery until all conditions met or waived</li> <li><input type="checkbox"/> Medical assessment completed by Hip and Knee Replacement Clinic designated physician unless referring authorized practitioner advises clinic (s)he will be responsible (see Relevant Assessments below)</li> <li><input type="checkbox"/> Medical clearance to indicate conditions to be treated, recommended treatment, and ASA score (see right)</li> <li><input type="checkbox"/> Surgical Pts assessed by consultants at Surgeon’s discretion, or designate, or as determined by screening criteria for clearance (see Consults section, below, for criteria)                             <ul style="list-style-type: none"> <li>• At minimum, review criteria in IM and Anaesthesia categories</li> <li>• Consider specialized wellness programs (Wt loss, tobacco cessation, community exercise programs) as required</li> <li>• Review exclusion criteria for day surgery Pt (refer to Exclusion Criteria for Day Surgery Candidacy)</li> </ul> </li> <li><input type="checkbox"/> After consultation with specialists for medical clearance and optimization the Pt will be sent back to the referring authorized practitioner/Hip and Knee Replacement Clinic who indicated responsibility for medical clearance and oversight of optimization</li> <li><input type="checkbox"/> If referring authorized practitioner responsible, Hip and Knee Replacement Clinic CM to monitor progress</li> <li><input type="checkbox"/> Chronic opioid use pre-operatively is a negative indicator of post-operative outcome and attempts should be made to eliminate preoperative use of opioids. Preoperative opioid tapering should be initiated as early as possible to improve outcomes</li> </ul>	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="background-color: #0056b3; color: white;">Status</th> <th style="background-color: #0056b3; color: white;">Description</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;"><b>1</b></td> <td>Healthy Pt</td> </tr> <tr> <td style="text-align: center;"><b>2</b></td> <td>Mild systemic disease</td> </tr> <tr> <td style="text-align: center;"><b>3</b></td> <td>Severe systemic disease (not incapacitating)</td> </tr> <tr> <td style="text-align: center;"><b>4</b></td> <td>Severe systemic disease that is a constant threat to life</td> </tr> <tr> <td style="text-align: center;"><b>5</b></td> <td>Moribund, not expected to live 24 hours</td> </tr> <tr> <td style="text-align: center;"><b>E</b></td> <td>Emergency Procedure</td> </tr> </tbody> </table>	Status	Description	<b>1</b>	Healthy Pt	<b>2</b>	Mild systemic disease	<b>3</b>	Severe systemic disease (not incapacitating)	<b>4</b>	Severe systemic disease that is a constant threat to life	<b>5</b>	Moribund, not expected to live 24 hours	<b>E</b>	Emergency Procedure
Status	Description														
<b>1</b>	Healthy Pt														
<b>2</b>	Mild systemic disease														
<b>3</b>	Severe systemic disease (not incapacitating)														
<b>4</b>	Severe systemic disease that is a constant threat to life														
<b>5</b>	Moribund, not expected to live 24 hours														
<b>E</b>	Emergency Procedure														
Optimization Wrap Up, Hand Over to Surgical Site and Discharge Planning															
<ul style="list-style-type: none"> <li><input type="checkbox"/> Pt cleared for surgery minimum 2 weeks prior to surgery date</li> <li><input type="checkbox"/> Pts requiring an extensive discharge plan will return for an extra visit to Hip and Knee Replacement Clinic to discuss plan</li> <li><input type="checkbox"/> CM to complete surgical Pt agreement once all consults, including medical clearance, have been completed and report(s) forwarded                             <ul style="list-style-type: none"> <li>• All surgical Pts to review their agreement with their CM and sign-off</li> <li>• CM must also sign-off</li> <li>• Pt variances from agreement/care path (e.g. DOS) communicated by CM to acute care sites</li> </ul> </li> <li><input type="checkbox"/> OR booking package completed, as required, and forwarded to appropriate AHS/Covenant Health sites upon surgical agreement completion (Follow on to Same Day Admit and Intra-Operative Surgery Sections, see below).</li> </ul>															

### Suggested Tools

- [Surgical Patient Agreement](#)



Relevant Assessments


<b>Pulmonary Embolism</b>	<b>Major Bleeding</b>
<ul style="list-style-type: none"> <li>☐ Assess pre-operatively for elevated risk (greater than standard risk). The following Pts are examples of those considered to be at elevated risk:             <ul style="list-style-type: none"> <li style="width: 50%;">• Previous documented pulmonary embolism</li> <li style="width: 50%;">• Previous history of Hypercoagulable states such as polycythemia</li> <li style="width: 50%;">• Previous history of thromboembolism</li> <li style="width: 50%;">• Spinal cord injury Pts</li> <li style="width: 50%;">• Previous history of cancer</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>☐ Assess pre-operatively for elevated risk (greater than standard risk). Pts with the following conditions are examples of those considered to be at elevated risk:             <ul style="list-style-type: none"> <li style="width: 50%;">• History of a bleeding disorder</li> <li style="width: 50%;">• History of recent hemorrhagic stroke</li> <li style="width: 50%;">• History of recent GI bleed</li> </ul> </li> </ul>
<b>OSA</b>	<b>PONV</b>
<ul style="list-style-type: none"> <li>☐ Assess pre-operatively for OSA:             <ul style="list-style-type: none"> <li>• Conduct Stop Bang test</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>☐ Assess pre-operatively for baseline predictive risk factors using the Apfel Simplified Risk Score:             <ul style="list-style-type: none"> <li style="width: 50%;">• Female Gender</li> <li style="width: 50%;">• History of PONV or motion sickness</li> <li style="width: 50%;">• Non-smoker</li> <li style="width: 50%;">• Intra-operative or post-operative use of opioids</li> </ul> </li> </ul>

**Anemia Management**

☐ Assess Pts pre-operatively for increased risk of low red blood cell mass using the AHS Preoperative Anemia Management & Hemoglobin Optimization Tool (2016).

**At-Risk Patient Populations:** Hgb <130g/L (male or female), weight <65kg, female gender, complex or revision surgery, renal disease, anti-platelet and/or anti-coagulant therapy, hematologic conditions (i.e.: Thalassemia), 'No Blood'/transfusion-refusal

**Ideal Timeline for Assessment:** Ideally at surgical INTAKE, at time of acceptance for surgery; **at least 30 days preop**



**Labs/Imaging**

<b>Standards:</b>	
<ul style="list-style-type: none"> <li>☐ All Pts tested per lab and ECG requirements (see table below for guidelines)</li> <li>☐ Pt-specific testing to monitor and achieve medical threshold defined in surgical Pt agreement; e.g. chest x-ray; INR for Pts on warfarin</li> <li>☐ Laboratory investigations should be ordered only when indicated by the Pt's medical status, drug therapy, or the nature of the proposed procedure</li> </ul>	<ul style="list-style-type: none"> <li>☐ ECGs are valid for three months, if available to anaesthesiologist, and if there have been no changes in symptoms in that time</li> <li>☐ Lab work is valid for 14 weeks</li> <li>☐ C-Spines are valid for one year</li> <li>☐ Chest x-rays are valid for one year</li> </ul>

**Table: Preoperative Laboratory Testing Guidelines for Common Comorbidities**

Note: This is minimum suggested pre-operative screening, tailor to the Pt's requirements for medical clearance

- Pre-operative **HbA1c, ferritin, and/or albumin** testing at Surgeon's discretion.
- **β-HCG** can be offered to premenopausal women who may be pregnant. Surgery need not be cancelled if the Pt declines.
- **Type and Screen** not routinely required. May be ordered, at Surgeon's discretion, for revisions or bilaterals.

Condition	CBC	Electrolytes	Creatinine eGFR	INR ±PTT	A1C	ECG	C-Spines	Chest X-ray
≥60 years old	X		X			X		
Revision Pts	X							
Inflammatory disease	X							
Known or suspected anemia, malnutrition, bleeding disorder, or bone marrow suppression	X							
Cardiovascular Disease (IHD CHF, Valvular HD, Pacemaker)	X		X			X		X
Hypertension		X	X			X		
Chronic Lung Disease (COPD, Pulmonary Fibrosis), Smokers >20/day	X							X
Diabetes			X		X	X		
Hepatic Disease (i.e. sclerosis)	X			X				
Renal Disease, Adrenocortical disease	X	X	X					
Therapy with diuretics, oral corticosteroids, lithium, DDAVP, or digoxin		X	X		X			
Pt taking Anticoagulants	X			X ±				
Malignancy	X							
Radiation or Chemotherapy in last 12 months	X							
Rheumatoid Arthritis							X	
Previous spinal instrumentation or fusion								
Immigrant without chest x-ray in last 12 months								X
± Coumadin = INR; No INR if therapy with warfarin stopped six days pre-operatively LMWH = No Action						Unfractionated heparin = PTT Antiplatelet Agents = No Action		

Consults for Consideration

Type of Consult	Criteria for Referral
<b>Internal Medicine</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Cardiovascular disease:               <ul style="list-style-type: none"> <li>• IHD with angina with mild exercise, or worsening angina</li> <li>• MI in last 12 months</li> <li>• Symptomatic valvular heart disease</li> <li>• CHF in last 12 months</li> <li>• Uncontrolled hypertension</li> </ul> </li> <li><input type="checkbox"/> Diabetic</li> <li><input type="checkbox"/> Neuromuscular Disease</li> <li><input type="checkbox"/> Pulmonary Disease:               <ul style="list-style-type: none"> <li>• Shortness of breath with mild exercise.</li> <li>• OSA</li> </ul> </li> <li><input type="checkbox"/> Restrictive lung disease</li> <li><input type="checkbox"/> Hematology:               <ul style="list-style-type: none"> <li>• Anticoagulant for any reason,</li> <li>• Anemia,</li> </ul> </li> <li><input type="checkbox"/> Morbid obesity (BMI ≥40),</li> <li><input type="checkbox"/> Pts currently being managed by an internist or cardiologist with an ASA 3 or greater score</li> </ul>
<b>Anaesthesia</b> (Also consider IM criteria for Anaesthesia referral)	<ul style="list-style-type: none"> <li><input type="checkbox"/> Known or suspected allergy to anesthetic drugs.</li> <li><input type="checkbox"/> Pt or family history of Malignant Hyperthermia</li> <li><input type="checkbox"/> Significant complications with previous anesthetics (including awareness).</li> <li><input type="checkbox"/> Jehovah's Witness – for hip arthroplasty</li> <li><input type="checkbox"/> History of Chronic Pain or long term (&gt;6month) Opioid Usage</li> <li><input type="checkbox"/> Pts currently being managed by an internist or cardiologist with an ASA 3 or greater score</li> </ul> <p>Note: AP lateral spine x-ray to be ordered and provided to anaesthetist for Pts with lumbar spine instrumentation or fusion</p> <p>Flexion extension cervical spine views to be ordered and provided to anaesthetist for Pts with rheumatoid arthritis</p> <ul style="list-style-type: none"> <li><input type="checkbox"/> Pt's request</li> <li><input type="checkbox"/> Known or suspected difficult airway</li> <li><input type="checkbox"/> Bleeding diathesis</li> <li><input type="checkbox"/> Morbid obesity (BMI ≥ 40)</li> </ul>
<b>Clinic OT Visit</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Pts with home issues and home visit not possible</li> <li><input type="checkbox"/> Out of region or major centre with limited access to OT resources</li> <li><input type="checkbox"/> Pt with significant ADL issues</li> <li><input type="checkbox"/> Require referral to AHS equipment program</li> </ul>
<b>Home Assessment/ Homecare Referral</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Pt having difficulty with self care, medication management or independent functioning pre-operatively</li> <li><input type="checkbox"/> Pt requiring extra (more permanent) equipment in home</li> <li><input type="checkbox"/> Pt not reliable historian</li> <li><input type="checkbox"/> Safety/fall risk</li> </ul>

Type of Consult	Criteria for Referral
<b>Nutritional Consult</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Pt with BMI greater than 30 or less than 18</li> <li><input type="checkbox"/> Anemic Pts</li> <li><input type="checkbox"/> Diabetic or hypertensive Pts with nutritional issues</li> <li><input type="checkbox"/> Pts with Chronic Kidney Disease</li> </ul>
<b>Group Pre-Operative Physiotherapy Treatments</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Frail elderly or debilitated Pt with pain and/or weakness limiting function at home pre-operatively</li> <li><input type="checkbox"/> Deconditioned Pt with poor upper extremity strength and/or ROM</li> <li><input type="checkbox"/> Deconditioned Pt with poor cardiovascular fitness and minimal exercise tolerance</li> <li><input type="checkbox"/> Pt with multiple joint involvement that limits function pre-operatively</li> <li><input type="checkbox"/> Pt with decreased balance and poor/unsafe ambulation pre-operatively</li> <li><input type="checkbox"/> Pt with significant contractures or quad lag pre-operatively</li> </ul>
<b>Individual Pre-Operative Physiotherapy Treatments</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Frail elderly or debilitated Pt with pain/and or weakness limiting function at home pre-operatively</li> <li><input type="checkbox"/> Pt has communication issues</li> <li><input type="checkbox"/> Pt has complex functional or medical issues</li> <li><input type="checkbox"/> Difficult or disruptive Pt</li> <li><input type="checkbox"/> Out of region or major centre with limited access to rehabbing treatment</li> </ul>
<b>Social Work</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Financial concerns</li> <li><input type="checkbox"/> Employment concerns</li> <li><input type="checkbox"/> Limited support system</li> <li><input type="checkbox"/> Difficult family dynamics</li> <li><input type="checkbox"/> Transportation issues</li> <li><input type="checkbox"/> Accommodation issues</li> <li><input type="checkbox"/> Addiction issues</li> <li><input type="checkbox"/> Alleged abuse</li> <li><input type="checkbox"/> Childcare issues</li> <li><input type="checkbox"/> Cultural/Language barriers</li> <li><input type="checkbox"/> Mental health or behavioral issues</li> <li><input type="checkbox"/> Concurrent crisis</li> <li><input type="checkbox"/> Legal issues</li> </ul>

Teaching

- Teaching is an imperative part of the entire Pt experience at the Hip and Knee Replacement Clinic

The Education Session	Equipment and Supplies
<ul style="list-style-type: none"> <li>□ <b>Format:</b> <ul style="list-style-type: none"> <li>• Site specific format to be determined at clinic team’s discretion</li> <li>• Education session completed by CM and others e.g. OT, PT</li> <li>• Consider grouping patients together based on day surgery candidacy or surgical approach to streamline delivery of customized messages.                             <ul style="list-style-type: none"> <li>○ Example: the importance of the stair climbing milestone for timely discharge</li> </ul> </li> </ul> </li> <li>□ <b>Key Items to Teach/Enforce:</b> <ul style="list-style-type: none"> <li>• Daily bath or shower with 4% chlorhexidine sponges for 3 days leading up to surgery</li> <li>• Topical application of 2% mupirocin* ointment twice per day to the nares for 5 days leading up to surgery<sup>4,5</sup></li> <li>• No lotion or cream to be used on the surgical site 5 days prior to surgery</li> <li>• No hair removal to be done 2 weeks prior to admission<sup>6</sup></li> <li>• Realistic post-operative pain expectations and functional goals</li> <li>• Pain management strategies, including non-drug strategies</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>□ Surgery Pt Teaching Book explaining aspects of intervention from beginning to end plus Phase I &amp; II exercises</li> <li>□ Classroom with comfortable (tall) chairs and tables for Pt to sit and write, bed to demonstrate transfers</li> <li>□ VCR, DVD and TV</li> <li>□ Telehealth or other virtual options</li> <li>□ Pt skin wash package including 4% chlorhexidine sponges</li> <li>□ 2% mupirocin ointment prescription</li> <li>□ Hip kit (reacher, long handled shoehorn/ stocking aid/ long handled bath sponge)</li> <li>□ Teaching crutches, walker, bathroom, stairs and dressing aids</li> <li>□ Theraband (exercise elastic) and exercise instructions</li> <li>□ Equipment list for Pts to organize for discharge (friend or family, vendors, RX or STELP, Health Unit)</li> <li>□ Available resources for Pts (Home Care, Meals on Wheels, Lifeline, etc.); Education Video/DVD</li> </ul>
<p><b>Activity / Mobility Instructions</b></p> <ul style="list-style-type: none"> <li>□ Home pre-operative strengthening program/exercises for all surgical Pts</li> <li>□ Use of walking aids as required</li> </ul>	<p><b>Surgical Agreement Completion</b></p> <ul style="list-style-type: none"> <li>□ All surgical Pts to review their agreement with their CM</li> <li>□ CM must arrange to have appropriate sign-offs completed before handoff to surgical site.</li> </ul>
<p><b>Patient/Family/Buddy Responsibility</b></p> <ul style="list-style-type: none"> <li>□ Buddy to attend             <ul style="list-style-type: none"> <li>• Surgical agreement discussion and signing</li> <li>• Pre-op teaching sessions and other appointments as necessary</li> </ul> </li> <li>□ Comply with pre-surgery optimization programs</li> <li>□ Prepare home and organize required equipment prior to surgical date</li> <li>□ Notify Hip and Knee Replacement CM if medical status changes</li> </ul>	<p><b>Suggested Tools</b></p> <ul style="list-style-type: none"> <li>📄 <a href="#">Minimally Invasive Approaches Handout</a></li> </ul>

\*This is an off-label use of this drug product; studies have associated it with very minimal risk to patient health and safety.<sup>4,5</sup>

## SECTION

### Same Day Admit to Intra-Operative Surgery

#### Same Day Admit (Pre-Surgery)

- Preparation
- Upon Admittance to Hospital
- Pre-Operative Medication Guidelines



#### Intra-Operative Surgery

- Surgical Guidelines
- Intra-Operative Medication Guidelines

**Same Day Admit (Pre-Surgery)**

**Preparation**

Patient/Family/Buddy Responsibility	Nutrition
<ul style="list-style-type: none"> <li><input type="checkbox"/> Bring all current medications</li> <li><input type="checkbox"/> Bring labeled reacher, dressing aids, exercise logbook, crutches and/or walker (Pt-specific aids)</li> <li><input type="checkbox"/> Buddy to accompany Pt to the hospital/site at scheduled time</li> <li><input type="checkbox"/> No hair removal to be done prior to admission<sup>6</sup></li> <li><input type="checkbox"/> Chlorhexidine skin washes night prior to or morning of surgery (sponge provided to Pt in Hip and Knee Replacement Clinic)</li> <li><input type="checkbox"/> Pts on Warfarin need an INR test the day before surgery, with an INR goal of <math>\leq 2</math></li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Follow Eating and Drinking Before Surgery Instructions:                             <ul style="list-style-type: none"> <li>• Eat as usual until 8 hours before surgery</li> <li>• May have a final light, low-fat snack before stopping all solids, 6 hours before surgery</li> <li>• Clear fluids only until 3 hours before surgery</li> <li>• Nothing by mouth 3 hours before surgery</li> </ul> </li> </ul>
Activity/Mobility	
<ul style="list-style-type: none"> <li><input type="checkbox"/> As directed in Pt Instruction Guide</li> </ul>	

**Upon Admittance to Hospital**

Assessment/Monitoring	Communication to Patient
<ul style="list-style-type: none"> <li><input type="checkbox"/> Vital signs <span style="margin-left: 150px;"><input type="checkbox"/> Physical assessment</span></li> <li><input type="checkbox"/> Review of pre-operative medications and complete or update BPMH</li> <li><input type="checkbox"/> Utilize Pt warming device (e.g. Bair Hugger) for warming 30 minutes pre-operatively</li> <li><input type="checkbox"/> No hair removal is optimal<sup>6</sup> <ul style="list-style-type: none"> <li>• Perform hair removal as needed—must be done 2 hours prior to entering surgical suite<sup>6</sup></li> </ul> </li> <li><input type="checkbox"/> Anesthesia check in with Pt at pre-op area</li> <li><input type="checkbox"/> Initiate IV access and fluids</li> <li><input type="checkbox"/> Start medication as per Anaesthesiologist or Surgeon’s orders. Specific doses depend on Pt’s risk (e.g. GI, Cadiac). See medication guidelines below</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Explain the OR process to the Pt</li> </ul>
	Tests
	<ul style="list-style-type: none"> <li><input type="checkbox"/> Diabetics receiving therapy need a glucometer reading on the morning of surgery</li> <li><input type="checkbox"/> If not done the day before, Pts on Warfarin need an INR on the morning of surgery, with an INR goal of <math>\leq 2</math></li> </ul>

Same Day Admit Pre-Operative Medication

Medication Type	Instructions										
<b>Antiemetics</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Use Apfel Simplified Risk Score to recognize Pts who are more likely to experience PONV</li> <li><input type="checkbox"/> Pts identified as high risk for PONV should be treated prophylactically</li> </ul> <p>Apfel Simplified Risk Score for PONV in adults</p> <table border="1" data-bbox="411 407 814 597"> <thead> <tr> <th data-bbox="411 407 705 440">Risk Factors</th> <th data-bbox="711 407 814 440">Points</th> </tr> </thead> <tbody> <tr> <td data-bbox="411 444 705 477">Female Gender</td> <td data-bbox="711 444 814 477">1</td> </tr> <tr> <td data-bbox="411 482 705 514">Non-Smoker</td> <td data-bbox="711 482 814 514">1</td> </tr> <tr> <td data-bbox="411 519 705 552">History of PONV</td> <td data-bbox="711 519 814 552">1</td> </tr> <tr> <td data-bbox="411 557 705 589">Intra-Op/Post-Op Opioids</td> <td data-bbox="711 557 814 589">1</td> </tr> </tbody> </table> <ul style="list-style-type: none"> <li><input type="checkbox"/> If the Apfel score is <math>\geq 3</math> AND Pt has history of PONV not responsive to usual care AND Pt will receive general anesthesia, consider aprepitant.</li> </ul>	Risk Factors	Points	Female Gender	1	Non-Smoker	1	History of PONV	1	Intra-Op/Post-Op Opioids	1
Risk Factors	Points										
Female Gender	1										
Non-Smoker	1										
History of PONV	1										
Intra-Op/Post-Op Opioids	1										
<b>Antibiotics<sup>7</sup></b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Provide dosing coverage for 24 hours post-operative to all patients</li> </ul>										
<b>Anti-Reflux</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Specific drug doses to be determined by Anesthesiologist or Surgeon depending on Pt's risk (e.g. GI, Cardiac)</li> </ul>										
<b>Analgesics</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Use multi-modal prophylaxis analgesia to control pain early</li> <li><input type="checkbox"/> Examples: NSAIDs, acetaminophen</li> </ul>										



Intra-operative – Surgery

Guidelines	Intra-operative Tests/Diagnostics
<ul style="list-style-type: none"> <li>□ <b>Start:</b> <ul style="list-style-type: none"> <li>• All cases start on time per schedule</li> <li>• Safe Surgery Checklists – all relevant staff to attend surgical-briefing<sup>8</sup></li> <li>• Site preparation with 2% CHG and 70% IPA (2%CHG-70%IPA) (first choice)<sup>9</sup>, if 2% CHG-70%IPA contraindicated, povidone/iodine (second choice) prep, 60% alcohol, and use iodine impregnated adhesive (Ioban) drape</li> <li>• Tourniquets used at Surgeon’s discretion</li> <li>• No routine use of Foley catheters<sup>10</sup></li> <li>• Surgeon signs site of incision and cuts through signature in the OR<sup>11</sup></li> </ul> </li> <li>□ <b>During Surgery:</b> <ul style="list-style-type: none"> <li>• Nursing assessments and monitoring per AHS/Covenant Health site policy</li> <li>• Pulse lavage to be available for use at Surgeon’s discretion -but no antibiotics</li> <li>• Safe Surgery Checklists – all relevant staff to attend surgical-time-out<sup>6</sup></li> <li>• Infiltration of joint with local anesthetic at Surgeon’s discretion</li> <li>• Minimum OR theatre temperature of 20-23°C <sup>6</sup></li> <li>• Utilize Pt warming device (e.g. Bair Hugger)</li> </ul> </li> <li>□ <b>Close:</b> <ul style="list-style-type: none"> <li>• No Hemovac drains for Hips or Knees<sup>12</sup></li> <li>• Safe Surgery Checklists – all relevant staff to attend surgical-debriefing<sup>8</sup></li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>□ Dependent upon Pt need and Surgeon’s discretion</li> </ul>
	Performance Standards
	<ul style="list-style-type: none"> <li>□ Where hospital sites are able: Pt’s surgery completed with dedicated team assigned to arthroplasty</li> <li>□ Benchmark duration for all primary elective procedures, including total hip, hip resurfacing, total knee and partial knee, is 75 minutes from incision to closure</li> <li>□ OR turnaround from closure on to incision next Pt &lt;45 minutes</li> <li>□ If a Surgeon cannot perform to the standard of 4 cases in a 7.5-hour day then OR day may be extended by up to 1.5 hours upon authorization by Surgical Chief and AHS/Covenant Health operations</li> </ul>

Intra-Operative Medication

Medication Type	Instructions																		
<b>Antiemetics</b>	<input type="checkbox"/> Apfel Simplified risk score for PONV in adults <table border="1" style="display: inline-table; vertical-align: middle;"> <thead> <tr> <th>Risk Factors</th> <th>Female Gender</th> <th>Non-Smoker</th> <th>History of PONV</th> <th>Post-Op Opioids</th> </tr> </thead> <tbody> <tr> <td>Points</td> <td>1</td> <td>1</td> <td>1</td> <td>1</td> </tr> </tbody> </table>	Risk Factors	Female Gender	Non-Smoker	History of PONV	Post-Op Opioids	Points	1	1	1	1								
	Risk Factors	Female Gender	Non-Smoker	History of PONV	Post-Op Opioids														
	Points	1	1	1	1														
	The following suggested selection of antiemetic can be titrated to clinical circumstances.																		
	<table border="1"> <thead> <tr> <th># Risk Factors</th> <th>Severity of PONV</th> <th>Prophylactic Strategy</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>10% Low</td> <td>No Prophylaxis</td> </tr> <tr> <td>1</td> <td>20% Low</td> <td>Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia</td> </tr> <tr> <td>2</td> <td>40% Moderate</td> <td>Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia +/- 5-HT3 antagonist (ondansetron 4 mg IV – can be given pre-op &amp; upon closure)</td> </tr> <tr> <td>3</td> <td>60% Severe</td> <td>Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia +/- 5-HT3 antagonist (ondansetron 4 mg IV – can be given pre-op &amp; upon closure) +/- NK1 antagonist (aprepitant 80 mg – given in Same Day Admit – see above)</td> </tr> <tr> <td>4</td> <td>80% Very Severe</td> <td>Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia +/- 5-HT3 antagonist (ondansetron 4 mg IV – can be given pre-op &amp; upon closure) +/- NK1 antagonist (aprepitant 80 mg – given in Same Day Admit – see above)</td> </tr> </tbody> </table>	# Risk Factors	Severity of PONV	Prophylactic Strategy	0	10% Low	No Prophylaxis	1	20% Low	Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia	2	40% Moderate	Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia +/- 5-HT3 antagonist (ondansetron 4 mg IV – can be given pre-op & upon closure)	3	60% Severe	Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia +/- 5-HT3 antagonist (ondansetron 4 mg IV – can be given pre-op & upon closure) +/- NK1 antagonist (aprepitant 80 mg – given in Same Day Admit – see above)	4	80% Very Severe	Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia +/- 5-HT3 antagonist (ondansetron 4 mg IV – can be given pre-op & upon closure) +/- NK1 antagonist (aprepitant 80 mg – given in Same Day Admit – see above)
	# Risk Factors	Severity of PONV	Prophylactic Strategy																
	0	10% Low	No Prophylaxis																
1	20% Low	Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia																	
2	40% Moderate	Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia +/- 5-HT3 antagonist (ondansetron 4 mg IV – can be given pre-op & upon closure)																	
3	60% Severe	Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia +/- 5-HT3 antagonist (ondansetron 4 mg IV – can be given pre-op & upon closure) +/- NK1 antagonist (aprepitant 80 mg – given in Same Day Admit – see above)																	
4	80% Very Severe	Steroid (dexamethasone 4 mg IV x 1 dose) at induction of anesthesia +/- 5-HT3 antagonist (ondansetron 4 mg IV – can be given pre-op & upon closure) +/- NK1 antagonist (aprepitant 80 mg – given in Same Day Admit – see above)																	
Note: Dexamethasone dose may be increased based on analgesic requirements – see below.																			
<b>Analgesics</b>	<input type="checkbox"/> Blocks – target is 90% spinal nerve blocks: <ol style="list-style-type: none"> <li>Consideration should be given to nerve blocks, particularly for Pts receiving chronic opioid therapy, or who have other complex pain histories:               <ul style="list-style-type: none"> <li>TKA: Femoral nerve block Or Adductor nerve block</li> </ul> </li> <li>Consideration should be given to intra-articular injections:               <ul style="list-style-type: none"> <li>TKA: 40 cc bupivacaine 0.25% or ropivacaine 0.5%, ketorolac 30 mg, and normal saline to a total volume of 150 cc</li> <li>THA: 40 cc bupivacaine 0.25% or ropivacaine 0.5%, ketorolac 30 mg, and normal saline to a total volume of 50 cc                   <ul style="list-style-type: none"> <li>Bupivacaine max dose should be 2 mg/kg</li> <li>Need to check GFR before giving ketorolac – avoid using if GFR &lt; 40</li> <li>0.5% ropivacaine should be used if there will be articular cartilage remaining</li> </ul> </li> </ul> </li> <li>Dexamethasone 0.1-0.25 mg/kg IV (max dose 20mg from both antiemetics and analgesic approach)</li> </ol> <input type="checkbox"/> <b>For Revision:</b> Pts may require additional epidural or general due to length of case																		

**Table: TXA Guidelines<sup>13</sup>**


Communication		Contraindications	
<ul style="list-style-type: none"> <li>□ Use of IV versus topical administration is at the Surgeon and Anaesthesiologist's discretion</li> <li>□ The plan for TXA and VTE prophylaxis must be discussed by the Surgeon and anesthesiologist at the start of the case</li> </ul>		<ul style="list-style-type: none"> <li>□ <b>General:</b> <ul style="list-style-type: none"> <li>• Allergy</li> <li>• Hypersensitivity to TXA</li> </ul> </li> <li>□ <b>Precautions</b> (Contraindications to IV use – consider Topical use instead):                             <ul style="list-style-type: none"> <li>• Pts at elevated risk of arterial or venous thrombosis                                     <ul style="list-style-type: none"> <li>▪ Within 3 months or Recurrent: acute DVT/PE</li> <li>▪ Within 12 months: prosthetic cardiac valve or drug-eluting stent and receiving clopidogrel, prasugrel or ticagrelor</li> <li>▪ Any anticoagulant therapy: e.g.: warfarin, DOAC, heparin, LMWH, etc.</li> <li>▪ Subarachnoid hemorrhage (potential for cerebral edema/infarction when given IV)</li> <li>▪ A-fib, A-flutter (no reliable safety data)</li> <li>▪ Received PCC or rFVIIa in past 4 hours</li> </ul> </li> <li>• Gross hematuria (potential ureter thrombosis)</li> <li>• Uncontrolled seizure disorder</li> <li>• Acquired disturbance of colour vision (prohibits assessment of one measure of toxicity)</li> <li>• Severe Hepatic or Renal disease (e.g.: Creatinine clearance &lt;30mLs/min)</li> </ul> </li> </ul>	
IV Dosing – Administered by Anaesthesiologist			
<ul style="list-style-type: none"> <li>□ Maximum infusion rate of 100 mg/min to avoid hypotension</li> </ul>			
TKA (tourniquet)	1 g IV infusion before tourniquet inflation <u>AND</u> 1 g IV infusion at tourniquet release		
TKA (NO tourniquet)	1-2 g IV infusion before incision		
THA	1-2 g IV infusion (OR 10-20 mg/kg) before incision		
Topical Dosing – Administered by Surgeon			
TKA & THA	2 g - 3 g in 50 mL - 100 mL NS Apply topically to joint for at least 3 minutes prior to closure		

## SECTION

### Acute Post-Operative Care

#### Post Operative Surgery Day 0-3

- Care to address each day patient is inpatient
- Acute Care Medication Guidelines



#### Discharge

- Care to accomplish prior to discharge
- Discharge to Recovery Medication Guidelines
- Discharge Referrals Criteria

**Post Operative Days 0-3**

Assessments/Monitoring/Interventions	Tests and Diagnostics
<ul style="list-style-type: none"> <li><input type="checkbox"/> <b>Follow prescriber's orders</b></li> <li><input type="checkbox"/> <b>Medical management</b> conducted by designated prescriber:               <ul style="list-style-type: none"> <li>• Designated prescriber may be: Surgeon, PCP, IM, Hospitalist or Resident-site specific</li> <li>• Review Pt history and pre-op medications on return from surgery</li> <li>• Designated prescriber to follow Acute Care Medication section (see below)</li> </ul> </li> <li><input type="checkbox"/> <b>Systems assessment</b> as per hospital protocol:               <ul style="list-style-type: none"> <li>• Skin assessment (Braden Scale) <span style="margin-left: 150px;">• Vital signs</span></li> <li>• Peripheral neurovascular assessment <span style="margin-left: 100px;">• Pain assessment</span></li> <li>• DB &amp; C q1h</li> <li>• Keep O2 sat greater than 92% or as prescribed</li> <li>• Assist Pt as needed with turning and positioning every 2 hours</li> <li>• Mechanical thromboprophylaxis at prescriber's discretion<sup>14</sup></li> <li>• If Neuroaxial anesthetic was used, assess for adequate motor function prior to mobilization of Pt</li> <li>• Fluid balance monitoring (IV, oral, urine)</li> <li>• Saline Lock IV when intake adequate; Discontinue IV upon D/C or when no longer clinically indicated</li> <li>• If Foley catheter required discontinue use early post-op day one</li> <li>• Maintain normothermia (as per Safer Healthcare Now)<sup>6</sup></li> <li>• Management of post-op hyperglycemia in diabetics (as per AHS Guidelines)<sup>15</sup></li> </ul> </li> <li><input type="checkbox"/> Assess dressing and provide interventions (as per prescriber's orders and site-specific Wound Care Guidelines)</li> <li><input type="checkbox"/> Intermittent cold therapy for knees<sup>16</sup></li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> For Pts symptomatic of low Hgb, do Hgb level and follow guidelines</li> </ul> <div style="border: 1px solid black; padding: 5px; margin: 5px 0;"> <p><b>Blood Transfusion Guidelines:</b></p> <p><b>Hgb &lt; 100g/l and</b> signs and symptoms of impaired O<sub>2</sub> delivery, heart rate ≥ 100, SBP ≤ 90, RR ≥ 20, Dyspnea, Syncope, Angina, Confusion, ECG ischemic changes  <b>Action:</b> Give O<sub>2</sub>, transfuse packed red blood cells 1 unit at a time and reassess</p> <p><b>Hgb ≥ 70 g/l</b> and no sign of impaired O<sub>2</sub> delivery, and no sign of cardiac history  <b>Action:</b> Monitor</p> <p><b>Hgb &lt; 70 g/l</b>  <b>Action:</b> Transfuse red blood cells sufficient to raise Hgb to greater than &gt; 70 g/l and reassess (1 unit should raise Hgb approximately 12 g/l)</p> </div> <ul style="list-style-type: none"> <li><input type="checkbox"/> CBC, Electrolytes, Creatinine, Urea POD 1. Repeat as needed</li> <li><input type="checkbox"/> If on Warfarin prior to surgery, daily PT (INR) with goal to resume therapeutic INR levels</li> <li><input type="checkbox"/> Glucose monitoring in diabetics as per post-op unit routine &amp; AHS Guidelines<sup>15</sup></li> <li><input type="checkbox"/> Any Pt-specific tests</li> <li><input type="checkbox"/> <b>One</b> post-operative x-ray required <u>within 12</u> weeks of surgery            Note: Timing is at Surgeon's discretion, may be completed during inpatient stay           <ul style="list-style-type: none"> <li>• <b>Hips:</b> <ul style="list-style-type: none"> <li>▪ AP pelvis centre 2"</li> <li>▪ Shoot through lateral affected hip to include stem</li> </ul> </li> <li>• <b>Knees:</b> AP and lateral</li> </ul> </li> </ul>
	<p><b>Supporting Tools</b></p> <p><a href="#">📄 Total Hip / Knee Arthroplasty Post-Operative Orders</a></p>

In-Hospital Consults		Nutrition
<input type="checkbox"/> PT <input type="checkbox"/> OT <input type="checkbox"/> Consults as required: <ul style="list-style-type: none"> <li>• Anaesthesia</li> <li>• IM</li> <li>• Cardiology</li> <li>• Pain Service</li> <li>• Dietitian</li> <li>• Others...</li> </ul>		<input type="checkbox"/> DAT - High Fibre (diet restrictions as ordered or in place pre-operatively)
Activity/Mobility		
<input type="checkbox"/> Rehabbing 2 x per day <ul style="list-style-type: none"> <li>• Independent ROM exercises between rehabbing visits</li> </ul> <input type="checkbox"/> Wt bearing as tolerated. <ul style="list-style-type: none"> <li>• <b>For Revision:</b> Activity and WB at Surgeon's discretion</li> </ul> <input type="checkbox"/> <b>Mobilization</b> to begin within 4 hours post-op <ul style="list-style-type: none"> <li>• Pts mobilized 10 steps or more on day of surgery</li> <li>• Encourage F/A exercises q1h</li> <li>• Transfers in/out of bed (assisted as required)</li> <li>• Up in chair for meals</li> </ul> <input type="checkbox"/> ADL practice with adaptive equipment as required		<input type="checkbox"/> For Hips ensure raised toilet seat/commode is in bathroom <input type="checkbox"/> <b>Precautions:</b> At Surgeon's discretion <input type="checkbox"/> Progressions Should be Observed: <ul style="list-style-type: none"> <li>• Towards independent bed/chair transfers</li> <li>• Walking in room, bathroom and hallway as able (minimum 3 - 5 x per day)</li> <li>• Increasing distance, using walker or crutches (assisted as required)</li> <li>• To crutches as able</li> <li>• To walk on stairs</li> </ul>
Patient/Family/Buddy Responsibilities		Teaching for Discharge Preparation
<input type="checkbox"/> Adhere to inpatient plan <input type="checkbox"/> Assist with ADLs <input type="checkbox"/> Family/buddy fill D/C prescription day before D/C <input type="checkbox"/> Ensure Pt has needed equipment <input type="checkbox"/> Ensure all supports are in place in preparation		<input type="checkbox"/> Reinforced precautions <input type="checkbox"/> Encourage Pt to record exercises in log book <input type="checkbox"/> Teach correct transfer techniques (bed/chair) <input type="checkbox"/> Ambulation/ROM instruction <input type="checkbox"/> Confirm home support services per surgical Pt agreement, or if required (See Discharge section for home care criteria)

Acute Care Medication

Medication Type	Instructions																							
<b>Bowel Management</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Assess and initiate bowel management</li> <li>Prevention of post-operative ileus with routine dosing of laxatives</li> </ul>																							
<b>Antiemetics</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> High-risk Pts will require an additional antiemetic to obtain adequate prophylaxis to prevent PONV</li> <li><input type="checkbox"/> Multimodal approach for prevention and treatment of PONV is recommended</li> <li>If Pt experiences PONV, despite adequate prophylaxis, medications from a different class must be used</li> </ul>																							
<b>Antibiotics<sup>7</sup></b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Provide dosing coverage for 24 hours post-operative to all patients</li> <li><input type="checkbox"/> Use same antibiotic as pre-op</li> </ul>																							
<b>Anticoagulants</b> <small>14, 17</small> see VTE Guidelines	<ul style="list-style-type: none"> <li><input type="checkbox"/> It is preferable to continue the same anticoagulant drug from pre-operative to post-operative</li> </ul> <p><b><i>American Academy of Orthopaedic Surgeons Clinical Guideline on Prevention of Pulmonary Embolism in Pts Undergoing Total Hip and Knee Arthroplasty (Modified to include Approved factor Xa inhibitors at recommended dose)</i></b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th colspan="3">Standard risk of Pulmonary Embolism</th> <th colspan="2">Elevated risk of Pulmonary Embolism</th> </tr> </thead> <tbody> <tr> <td rowspan="2"><b>Standard risk for major bleeding</b></td> <td>Aspirin</td> <td colspan="2">Synthetic Pentasaccharides</td> <td>LMWH</td> <td>Synthetic Pentasaccharides</td> </tr> <tr> <td>Direct Factor Xa Inhibitor</td> <td>Warfarin</td> <td>LMWH</td> <td>Direct Factor Xa Inhibitor</td> <td>Warfarin</td> </tr> <tr> <td><b>Elevated risk for major bleeding</b></td> <td>Aspirin</td> <td colspan="2">Warfarin</td> <td colspan="2">None</td> </tr> </tbody> </table>		Standard risk of Pulmonary Embolism			Elevated risk of Pulmonary Embolism		<b>Standard risk for major bleeding</b>	Aspirin	Synthetic Pentasaccharides		LMWH	Synthetic Pentasaccharides	Direct Factor Xa Inhibitor	Warfarin	LMWH	Direct Factor Xa Inhibitor	Warfarin	<b>Elevated risk for major bleeding</b>	Aspirin	Warfarin		None	
		Standard risk of Pulmonary Embolism			Elevated risk of Pulmonary Embolism																			
	<b>Standard risk for major bleeding</b>	Aspirin	Synthetic Pentasaccharides		LMWH	Synthetic Pentasaccharides																		
		Direct Factor Xa Inhibitor	Warfarin	LMWH	Direct Factor Xa Inhibitor	Warfarin																		
	<b>Elevated risk for major bleeding</b>	Aspirin	Warfarin		None																			
<b>Duration:</b>																								
<ul style="list-style-type: none"> <li><input type="checkbox"/> Duration should be 10-35 days from initiation of medication, at the Surgeon's discretion of patient tailored medication regime</li> </ul>																								

Medication Type	Instructions				
<b>Analgesics</b> <sup>18</sup>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Adequate pain relief post-operatively is essential to allow patients to fully participate in post-operative protocols and meet discharge goals. Attempt to maintain pain score less than 4/10</li> <li><input type="checkbox"/> Communication regarding pain management orders between pain management, pharmacy, nursing and orthopedic Surgeon to occur as needed</li> <li><input type="checkbox"/> Multimodal approach to pain control is optimal, employing non-opioid analgesics, nonsteroidal, opioids and nonpharmacological modalities</li> <li><input type="checkbox"/> Patients will typically require opioid medication post-op. Opioid medication should be titrated to the minimum dose that allows patients to fully participate in post-operative protocols and meet discharge goals</li> <li><input type="checkbox"/> Typical patients will require tapering doses of opioid medications for 6-12 weeks post-operative with the earliest discontinuation as possible being optimal</li> <li><input type="checkbox"/> Long acting opioids should be avoided, unless patients are already on long acting opioids in the community</li> </ul> <p>Individual Pt assessments to be considered for providing analgesia:</p>				
	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="background-color: #0056b3; color: white;">Drug Category</th> <td style="width: 33%;"> <b>Acetaminophen</b> (maximum 4g in 24 hours from all sources)         </td> <td style="width: 33%;"> <b>NSAIDs</b> </td> <td style="width: 33%;"> <b>Opioids</b> </td> </tr> </table>	Drug Category	<b>Acetaminophen</b> (maximum 4g in 24 hours from all sources)	<b>NSAIDs</b>	<b>Opioids</b>
	Drug Category	<b>Acetaminophen</b> (maximum 4g in 24 hours from all sources)	<b>NSAIDs</b>	<b>Opioids</b>	
	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="background-color: #0056b3; color: white;">Cautions</th> <td style="width: 33%;">Liver disease</td> <td style="width: 33%;">Creatinine clearance &lt;50</td> <td style="width: 33%;">Elderly – use lower dosing</td> </tr> </table>	Cautions	Liver disease	Creatinine clearance <50	Elderly – use lower dosing
	Cautions	Liver disease	Creatinine clearance <50	Elderly – use lower dosing	
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="background-color: #0056b3; color: white;">Cautions</th> <td style="width: 33%;">Elderly – use lower dosing</td> <td style="width: 33%;">History of GI bleed</td> <td style="width: 33%;">Opioid naïve – use lower dosing</td> </tr> </table>	Cautions	Elderly – use lower dosing	History of GI bleed	Opioid naïve – use lower dosing	
Cautions	Elderly – use lower dosing	History of GI bleed	Opioid naïve – use lower dosing		
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="background-color: #0056b3; color: white;">Cautions</th> <td style="width: 33%;">-</td> <td style="width: 33%;">Consider continuing Pts' regular pre-operative NSAIDs regime</td> <td style="width: 33%;">Consider individualized dosing in Pts on regular pre-operative opioid regime to account for increased dosing needs</td> </tr> </table>	Cautions	-	Consider continuing Pts' regular pre-operative NSAIDs regime	Consider individualized dosing in Pts on regular pre-operative opioid regime to account for increased dosing needs	
Cautions	-	Consider continuing Pts' regular pre-operative NSAIDs regime	Consider individualized dosing in Pts on regular pre-operative opioid regime to account for increased dosing needs		
<b>Pruritus</b>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Ensure supportive therapy medications are ordered</li> </ul>				



Discharge

Criteria for Patient Discharge – All Must Be Met	Suggested Tools
<ul style="list-style-type: none"> <li><input type="checkbox"/> Eating and elimination patterns are approximating, or have returned to, normal</li> <li><input type="checkbox"/> Surgical incision is clean and dry, or arrangements for wound care have been made</li> <li><input type="checkbox"/> Pt is able to:                             <ul style="list-style-type: none"> <li>• Do bed and chair transfers independently</li> <li>• Demonstrate movement precautions</li> <li>• Perform ADL and use necessary equipment and aids</li> <li>• Walk independently with appropriate aids on the level and stairs – maintaining any WB restriction</li> </ul> </li> </ul> <p><b>Note: Pts may be discharged when not independent with any of above, <u>if appropriate assistance is available</u></b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> Appropriate D/C destination, transportation, equipment and any services (i.e. referrals) required are confirmed to be in place</li> <li><input type="checkbox"/> D/C prescriptions ordered by designated prescriber (see Recovery Medication section below) Note: additional/different medications required due to surgical or inpatient events to be determined by prescriber Note: BPMH reconciliation completed prior to D/C</li> <li><input type="checkbox"/> D/C teaching instructions and plan (documented in site-specific Pt D/C summary form) have been <u>received and understood</u> by the Pt, buddy and/or family</li> </ul>	<ul style="list-style-type: none"> <li><a href="#">Patient Discharge Summary</a></li> <li><a href="#">Going Home After a Nerve Block</a></li> </ul>
	Guidelines to Staff Achieving Discharge
	<ul style="list-style-type: none"> <li><input type="checkbox"/> Identify barriers to D/C early</li> <li><input type="checkbox"/> Identify potential variance from Pt agreement designed by Hip and Knee Clinic CM and <b>maintain regular communication with CM</b> <sup>19</sup></li> <li><input type="checkbox"/> Confirm responsibility for medical management post D/C</li> <li><input type="checkbox"/> PCP engaged as needed and provided with inpatient summary and/or D/C information</li> <li><input type="checkbox"/> Complete D/C information summary upon D/C</li> </ul> <p>Goal: D/C Pt without complications, as planned and scheduled</p> <p>Goal: D/C Pt home wherever possible</p>
	Referral Arrangement
	<ul style="list-style-type: none"> <li><input type="checkbox"/> Referrals arranged and confirmed as required:                             <ul style="list-style-type: none"> <li>• Homecare (see criteria below)</li> <li>• Outpatient PT (see criteria below)</li> <li>• Sub-acute transfer (see sub-acute section below)</li> </ul> </li> </ul>
Teaching Instructions and Discharge Plan Includes	Patient/Family/Buddy Responsibility
<ul style="list-style-type: none"> <li><input type="checkbox"/> Topics to be covered by Nurse/PT/OT include:                             <ul style="list-style-type: none"> <li>• Follow-up appointments at Hip and Knee Clinic</li> <li>• Exercises and precautions</li> <li>• Referrals (see right)</li> <li>• Incision care</li> <li>• Options for support between D/C and 2-week follow-up, as per D/C summary sheet:                                     <ul style="list-style-type: none"> <li>○ When to call CM</li> <li>○ When to call GP</li> </ul> </li> </ul> </li> <li>• Medication:                             <ul style="list-style-type: none"> <li>▪ Required prescriptions procurement</li> <li>▪ Anticoagulant administration</li> <li>▪ Analgesic administration and pain management</li> <li>▪ Appropriate analgesia tapering</li> <li>▪ Safe opioid storage and disposal at a local pharmacist</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Ensure understanding of D/C teaching instructions</li> <li><input type="checkbox"/> Family/buddy attend D/C instruction review with Pt and inpatient staff</li> <li><input type="checkbox"/> D/C medication brought to inpatient unit for review and reconciliation prior to D/C</li> <li><input type="checkbox"/> Family/buddy transport Pt back home</li> </ul>

Discharge to Recovery Medication

Medication Type	Discharge to 2 Week Post-Op	2-6 Week Post-Op	6-12 Week Post-Op + Monitoring
<b>Bowel Management</b>	<input type="checkbox"/> Sennosides 8.6 mg oral prn (HOLD IF stool loose) <input type="checkbox"/> Glycerine suppository prn <input type="checkbox"/> PEG 3350 (polyethylene glycol 3350) 17 g in 250 mL fluid orally daily for constipation (HOLD IF stool loose)		N/A
<b>Antibiotics</b>	<input type="checkbox"/> Routine antibiotic prophylaxis is not indicated for dental Pts with total joint replacements. At Surgeon discretion for high risk Pts. <sup>20</sup>		
<b>Analgesics<sup>18</sup></b>	<input type="checkbox"/> Adequate pain relief post-operatively is essential to allow patients to fully participate in post-operative protocols and meet discharge goals. Attempt to maintain pain score less than 4/10 <input type="checkbox"/> Communication regarding pain management orders between pain management, pharmacy, nursing and orthopedic Surgeon to occur as needed <input type="checkbox"/> Multimodal approach to pain control is optimal, employing non-opioid analgesics, nonsteroidal, opioids and nonpharmacological modalities <input type="checkbox"/> Patients will typically require opioid medication post-op. Opioid medication should be titrated to the minimum dose that allows patients to fully participate in post-operative protocols and meet discharge goals <input type="checkbox"/> Typical patients will require tapering doses of opioid medications for 7 days (THA) and 14 days (TKA) post-operative with the earliest discontinuation as possible being optimal. Devise an individualized plan for tapering to avoid opioid dependence <input type="checkbox"/> Long-acting opioids should be avoided, unless patients are already on long acting opioids in the community <input type="checkbox"/> Chronic opioid use preoperatively is a negative indicator of post-operative outcome and attempts should be made to eliminate preoperative use of opioids		

Medication Type	Discharge to Community Instructions
<b>Anticoagulants<sup>14, 17</sup></b> see VTE Guidelines	<input type="checkbox"/> Follow anticoagulation drug established at D/C from acute care centre <input type="checkbox"/> Continue anticoagulation drug for 10-35 days, at Surgeon's discretion

Discharge Referrals for Consideration

Type of Referral	Criteria
<p><b>Homecare Post-Operative</b></p>	<ul style="list-style-type: none"> <li>□ Requires personal care assistance to be safe with certain ADL e.g. bathing, walking, bed mobility, dressing, feeding, off and on toilet</li> <li>□ Pt's condition has changed and a home visit is required to assess and recommend equipment or strategies that will improve Pt safety and independent function at home</li> <li>□ Unable to:               <ul style="list-style-type: none"> <li>• administer required medication</li> <li>• change a wound dressing (if required) and/or requires monitoring of a draining incision</li> <li>• leave the home for required physiotherapy treatments, exercises or monitoring</li> </ul> </li> </ul>
<p><b>Outpatient Physiotherapy Treatments Post-Operative</b></p>	<ul style="list-style-type: none"> <li>□ <b>TKA, Knee Revision or Partial Knee</b> (if treatment is required, usually at D/C to 6 weeks post-op)           <ul style="list-style-type: none"> <li>• Poor gait pattern or balance:               <ul style="list-style-type: none"> <li>a. If requires gait correction within WB restriction</li> <li>b. If requires progression of WB (if Pt is Wt bear as tolerated or has been given new WB orders)</li> </ul> </li> <li>• Poor quad contraction:               <ul style="list-style-type: none"> <li>a. &gt;15° quad lag</li> <li>b. &lt; Grade 2+ strength</li> <li>c. Unable to do a straight leg raise against gravity</li> </ul> </li> <li>• Pain and Swelling Control</li> <li>• Poor ROM:               <ul style="list-style-type: none"> <li>a. &lt;70° flexion</li> <li>b. &gt;15° flexion contracture</li> </ul> </li> <li>• If modalities decrease pain or swelling, making exercises more successful</li> <li>• If Pt:               <ul style="list-style-type: none"> <li>▪ would benefit from education regarding pain/swelling control techniques, activities and positions</li> <li>▪ needs instruction with walking aids (weaning off aid when ordered)</li> <li>▪ needs balance training or strengthening (to improve balance)</li> </ul> </li> </ul> </li> <li>□ <b>THA, Hip Resurfacing or Hip Revision</b> (if treatment is required, usually at D/C to 6 weeks post-op)           <ul style="list-style-type: none"> <li>• Poor hip ROM:               <ul style="list-style-type: none"> <li>a. &lt; 45° flexion, &lt;15° abduction</li> </ul> </li> <li>• Poor hip strength:               <ul style="list-style-type: none"> <li>a. &lt; Grade 2+ flexor strength</li> <li>b. &lt; Grade 2 abductor strength</li> </ul> </li> <li>• Significant edema in the surgical leg (pitting, causing decreased leg ROM)</li> <li>• If Pt would benefit from education regarding pain/swelling control techniques, activities and positions</li> <li>• Poor gait pattern or balance:               <ul style="list-style-type: none"> <li>a. If requires gait correction within WB restriction</li> <li>b. If requires progression of WB (if Pt is WB as tolerated or has been given new WB orders)</li> <li>c. If Pt needs instruction with walking aids (weaning down or off aid when ordered)</li> <li>d. If Pt needs balance training or strengthening (to improve balance)</li> </ul> </li> </ul> </li> </ul>

## SECTION

# Recovery<sup>21</sup>



**Discharge – 2 Weeks Post-Operative**

**Case Management Team**

- Complete follow up call with Pts within 24 hours of D/C
- Reinforce options for communication with CM and GP, as per Discharge Summary Instructions handout

**2 Weeks Post-Operative Visit <sup>21</sup>**

**Clinic Assessment**

- During Hip and Knee Replacement Clinic Team Evaluation within 10-14 days:**
  - Staples/suture removal, as required
  - Confirm surgical Pt agreement and plan adhered to
  - CM/Team assess incision condition, i.e. swelling, pain:
    - Infection assessment as per AAOS guidelines<sup>22</sup>
  - Rehabilitation/therapy assessment:
    - Assess ROM and gait
    - Confirm Pt maintaining precautions and WB status
    - Confirm using walking aid, bathroom equipment, dressing aids
    - Complete post-op Physiotherapy Referral/Report
  - Confirm medication as per Discharge to Recovery Medication section (see above) and Pt's D/C summary form
    - Address changes to individualized opioid tapering plan
    - Obtain altered prescriptions, as required
  - Communicate with Pt's referring authorized practitioner or PCP regarding Pt issues, as required
  - Confirm referral arrangement, as required

**Referral Arrangement**

- Home Care, OT or PT if required (See Discharge Referrals section, above)
- 7 group PT sessions are available for post-op rehab for Pts requiring functional optimization

**Tests / Diagnostics**

- If on Warfarin prior to surgery, complete daily PT (INR) post-operatively with goal to resume therapeutic INR levels (continue as ordered by GP/PCP)

**Nutrition**

- DAT – maintain well balanced diet or diet as specified re. Canada Food Guide

**Patient Goals for Activity/Mobility**

- Independent function:
  - Transfers
  - Dressing
  - Ambulation/stairs
  - Home exercise program
  - Self care
- Progressive return to normal daily activities, as tolerated
- Progressive walking distance, as tolerated

**Teaching**

- Educate Pt regarding pain medication, incision care, and potential complications (including swelling)

**Patient/Family/ Buddy Responsibility**

- Encourage independent exercise and mobilization
- Support at home as needed (laundry, driving, meal prep, etc.)

**6 Weeks and 12 Weeks Post-Operative Visit<sup>21</sup>**

<b>Clinic Assessment</b>	<b>Test / Diagnostics</b>
<ul style="list-style-type: none"> <li><input type="checkbox"/> <b>During Hip and Knee Replacement Clinic Team Evaluation at 6 and 12 weeks:</b> <ul style="list-style-type: none"> <li>• Pt is assessed by Surgeon and team, as appropriate                             <ul style="list-style-type: none"> <li>▪ Assess infection assessment as per AAOS guidelines<sup>22</sup></li> </ul> </li> <li>• Confirm medication as per Discharge to Recovery Medication section (see above) and Pt's D/C summary form                             <ul style="list-style-type: none"> <li>▪ Address changes to individualized opioid and foundational analgesia tapering</li> <li>▪ Obtain altered prescriptions, as required</li> </ul> </li> <li>• Communicate with Pt's referring authorized practitioner or PCP regarding Pt issues, as required</li> <li>• Referrals to post-operative out-Pt rehab, as required (see Discharge Referrals section, above)</li> </ul> </li> <li><input type="checkbox"/> <b>6 Week Visit Only:</b> <ul style="list-style-type: none"> <li>• Confirm removal of bathroom equipment and dressing aids, as able</li> <li>• Rehabilitation/therapy assessment, at Surgeon's discretion:                             <ul style="list-style-type: none"> <li>▪ Confirm WB, exercises, walking aids progressed and activities progressed</li> </ul> </li> </ul> </li> <li><input type="checkbox"/> <b>12 Week Visit Only:</b> <ul style="list-style-type: none"> <li>• Confirm Pt achieved outcomes as defined in Surgical Pt Agreement</li> <li>• Complete Hip/Knee Patient Reported Outcome and Experience Measures</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> INR for Pts on Warfarin (continue as ordered by GP/PCN)</li> <li><input type="checkbox"/> <b>One</b> post-operative x-ray required <u>within 12 weeks</u> of surgery Note: timing is at Surgeon's discretion, may be completed during in-Pt stay</li> <li>• <b>Hips:</b> <ul style="list-style-type: none"> <li>▪ AP pelvis centre 2" low</li> <li>▪ Lateral of affected hip</li> </ul> </li> <li>• <b>Knees:</b> <ul style="list-style-type: none"> <li>▪ AP, lateral and skyline view of affected knee</li> </ul> </li> </ul>
	<b>Patient/Family/ Buddy Responsibility</b>
	<ul style="list-style-type: none"> <li><input type="checkbox"/> Encourage independent exercise and mobilization</li> <li><input type="checkbox"/> Support at home as needed (laundry, driving, meal prep, etc.)</li> </ul>
	<b>Patient Goals for Activity/Mobility</b>
	<ul style="list-style-type: none"> <li><input type="checkbox"/> Independent function: Transfers, Ambulation/stairs, Dressing, Self-care</li> <li><input type="checkbox"/> Possible return to driving, at Surgeon's discretion</li> <li><input type="checkbox"/> Progressive walking distance, as tolerated</li> <li><input type="checkbox"/> <b>6 Weeks:</b> continue home exercise; <b>12 Weeks:</b> exercise in community, as advised</li> <li><input type="checkbox"/> <b>6 Weeks:</b> WB as tolerated, progress to cane if able, full WB if cleared by Surgeon</li> <li><input type="checkbox"/> Progress to Recovery Exercises, as directed at relevant visit:             <ul style="list-style-type: none"> <li>• Gravity resist, Theraband, light weights, ROM (gently into flexion past 90°)</li> </ul> </li> </ul>
<b>Patient Reported Outcome and Experience Measures</b>	
<ul style="list-style-type: none"> <li><input type="checkbox"/> At <b>6 Weeks</b> Pt completes:             <ul style="list-style-type: none"> <li>• Pt Feedback Survey</li> </ul> </li> <li><input type="checkbox"/> At <b>12 Weeks</b> Pt completes:             <ul style="list-style-type: none"> <li>• EQ5D-5L</li> <li>• WOMAC</li> </ul> </li> </ul>	
<b>Nutrition</b>	
<ul style="list-style-type: none"> <li><input type="checkbox"/> DAT – maintain well balanced diet or diet as specified re. Canada Food Guide</li> </ul>	
<b>Suggested Tools</b>	
<ul style="list-style-type: none"> <li><input type="checkbox"/> <a href="#">Hip and Knee Outcomes Tool (Combined WOMAC and EQ5D-5L)</a></li> <li><input type="checkbox"/> <a href="#">Hip and Knee Patient Feedback Survey</a></li> </ul>	
	<b>Teaching</b>
	<ul style="list-style-type: none"> <li><input type="checkbox"/> <b>At Surgeon's discretion:</b> Educate Pt regarding progression of exercises and functional movements</li> <li><input type="checkbox"/> Discuss safe opioid storage and disposal at local pharmacy</li> </ul>

## 1 Year Post-Operative Visit and Monitoring<sup>21</sup>

<p><b>1 Year Clinic Assessment</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> <b>During Hip and Knee Replacement Clinic Team Evaluation at 6 and 12 weeks:</b> <ul style="list-style-type: none"> <li>• Pt is assessed by Surgeon and team, as appropriate           <ul style="list-style-type: none"> <li>▪ Assess infection assessment as per AAOS guidelines<sup>22</sup></li> </ul> </li> <li>• Surgeon reviews all x-rays</li> <li>• Communicate with Pt's referring authorized practitioner or PCP regarding Pt issues, as required</li> <li>• Referrals to post-op out-Pt rehab, as required</li> <li>• Complete Hip/Knee Patient Reported Outcome Measures</li> </ul> </li> </ul>	<p><b>Suggested Tools</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> <a href="#">Hip and Knee Outcomes Tool (Combined WOMAC and EQ5D-5L)</a></li> </ul>
<p><b>Patient Reported Outcome Measures at 1 Year Visit</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> Pt completes:       <ul style="list-style-type: none"> <li>• EQ5D-5L</li> <li>• WOMAC</li> </ul> </li> </ul>	<p><b>Tests / Diagnostics</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> X-Rays at 1 year, and then aligned with recall frequency       <ul style="list-style-type: none"> <li>• <b>Hips:</b> <ul style="list-style-type: none"> <li>▪ AP pelvis centre 2" low</li> <li>▪ Shoot through lateral of affected hip</li> </ul> </li> <li>• <b>Knees:</b> <ul style="list-style-type: none"> <li>▪ AP and lateral of affected knee</li> <li>▪ Merchant view of patella</li> </ul> </li> </ul> </li> </ul>
<p><b>Monitoring Beyond 1 Year</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> Goal: 100% primary joint Pts monitored every 2 years</li> <li><input type="checkbox"/> <b>Primary surgeries:</b> monitoring is completed every five years, at Surgeon's discretion</li> <li><input type="checkbox"/> <b>Revision surgeries:</b> monitoring is completed annually, at Surgeon's discretion</li> <li><input type="checkbox"/> Monitoring Actions:       <ul style="list-style-type: none"> <li>• Surgeon's review of x-rays</li> <li>• In-clinic assessment</li> </ul> </li> <li><input type="checkbox"/> If prosthesis problem/issue/failure is identified, follow-up should be expedited<sup>21</sup></li> <li><input type="checkbox"/> Routine antibiotic prophylaxis is not indicated for dental Pts with total joint replacements. At Surgeon discretion for high risk Pts<sup>20</sup>.</li> </ul>	<p><b>Patient Goals for Activity/Mobility</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> Independent function</li> <li><input type="checkbox"/> Normal Daily activities</li> <li><input type="checkbox"/> Exercise in Community</li> </ul> <p><b>Patient/Family/ Buddy Responsibility</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> Encourage normal function</li> </ul> <p><b>Nutrition</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> DAT – maintain well balanced diet or diet as specified re. Canada Food Guide</li> </ul>

## SECTION

### Care of Patients in Subacute



## Subacute Care

### Criteria for Transferring to Subacute Care (Primary and Revision) May Include:

- Unable to manage environment at residence, e.g. no home support, difficult living arrangements (stairs, levels, access to bath/kitchen)
- Frail elderly with comorbidities
- Daily need for rehabilitative services and/or limited access to rehabilitation services
- Bilateral joint surgery
- Post-operative complications

### Information to Provide to Subacute Care Centre

- Inpatient D/C Information
- Relevant Precautions list/WB Status (prescribed at Surgeon's discretion)
- Include assessment of knee ROM (instructions to refer to community physio if less than 70° flexion and/or quad lag)
- Follow up appointment instructions (suture/staple removal) at 2 weeks' post-op
- General information sheet on THA/TKA for health care providers including
  - a. Brief description of surgery:
  - b. Copy of Surgical Pt Agreement (surgery – 1 year) so they understand plan:
    - Activity / Rehab expectations at subacute stage (including self care/ exercises/ activity)
    - Work toward independent transfers/ambulation/self care and dressing/ exercises
    - More time spent in activity than resting in bed (improving endurance, strength and function)
    - Issues to address for D/C from subacute facility to community
- Any nursing care required at home (dressing changes/ medication/bath)
- Any home adaptations and equipment required
- Any support/supervision required at home (family/friends/home care)
- Any meal/homemaking services needed at home
- Any exercise assistance/monitoring required at home
- Expectations/plan of activity/follow-up/outcome for community phase (subacute-12months)
- Contact phone numbers for questions/advice required by health care providers (clinic #)

### Medications

- |   |   |
|---|---|
| <ul style="list-style-type: none"> <li><input type="checkbox"/> <b>Bowel management:</b> see Acute Care Medication (page 23)</li> <li><input type="checkbox"/> <b>Antiemetics:</b> see Acute Care Medication (page 23)</li> <li><input type="checkbox"/> <b>Anticoagulants</b><sup>12, 15</sup>: see Acute Care Medication (page 23)</li> </ul> | <ul style="list-style-type: none"> <li><input type="checkbox"/> <b>Analgesics</b><sup>18</sup>:                     <ul style="list-style-type: none"> <li>• Check that analgesics/anti-inflammatories from acute care have been reassessed</li> <li>• See Acute Care Medication (page 23)</li> </ul> </li> </ul> |
|---|---|

Assessment / Monitoring for a Duration of Stay	Consults	Nutrition
<ul style="list-style-type: none"> <li><input type="checkbox"/> Per Surgeon hip or knee orders:               <ul style="list-style-type: none"> <li>• Adhere to standardized care path</li> <li>• Adhere to surgical Pt agreement and plan</li> <li>• Neurovascular and Physical Assessments</li> <li>• Dressing changes if needed</li> </ul> </li> <li><input type="checkbox"/> Communications:               <ul style="list-style-type: none"> <li>• Hip and Knee Clinic CM communicates with designated subacute contact re potential variance from surgical Pt agreement</li> <li>• CM or Pt's referring or PCP to oversee medical management</li> </ul> </li> <li><input type="checkbox"/> D/C:               <ul style="list-style-type: none"> <li>• Pt discharged without complications as planned and scheduled</li> <li>• Subacute Record completed upon D/C and faxed back to Hip and Knee Clinic</li> </ul> </li> </ul>	<input type="checkbox"/> As required	<input type="checkbox"/> DAT -High Fibre (Diet restrictions as ordered or in place pre-operatively)
<b>Activity/Mobility Goals</b>		
<ul style="list-style-type: none"> <li><input type="checkbox"/> Assist with a.m. care as required</li> <li><input type="checkbox"/> Rehabbing exercises 2 -3 x per day on unit</li> <li><input type="checkbox"/> Up to bathroom at night</li> <li><input type="checkbox"/> Up in chair for meals</li> <li><input type="checkbox"/> Progress mobilization to minimum of 5X per day with walker or crutches, maintaining WB restrictions</li> <li><input type="checkbox"/> Independent ROM exercises between rehabbing visits</li> <li><input type="checkbox"/> Progress independent function Transfers/ Ambulation/ Stairs/ Self care/ Dressing</li> <li><input type="checkbox"/> Pt to dress in own clothes (with/without assist)</li> </ul>		
Discharge Teaching	Patient / Family / Buddy Responsibility	
<ul style="list-style-type: none"> <li><input type="checkbox"/> Anticoagulant self administration taught and supervised</li> <li><input type="checkbox"/> Analgesic administration taught</li> <li><input type="checkbox"/> Home Exercises</li> <li><input type="checkbox"/> ADL instruction: dressing, tub transfers, car transfers</li> <li><input type="checkbox"/> Equipment reviewed (dressing aids to be used as required)</li> <li><input type="checkbox"/> D/C instructions reinforced/completed by nurse</li> <li><input type="checkbox"/> Confirm follow-up appointments</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Encourage independent exercise and mobilization</li> <li><input type="checkbox"/> Home prepared</li> <li><input type="checkbox"/> Support available</li> <li><input type="checkbox"/> Arrangements made for transport</li> <li><input type="checkbox"/> Transport home with goal of 10:00h</li> <li><input type="checkbox"/> D/C prescriptions filled by Pt support day before D/C and brought to inpatient unit for review and reconciliation prior to D/C.</li> </ul> <p>Note: additional/different medications required due to surgical or inpatient events to be determined by prescriber. Modified prescription(s) to be filled by Pt support</p>	

## References

1. College of Physicians & Surgeons of Alberta. Referral Consultation: Advice to the Profession. January 2017. [http://www.cpsa.ca/wp-content/uploads/2017/01/AP\\_Referral-Consultation.pdf](http://www.cpsa.ca/wp-content/uploads/2017/01/AP_Referral-Consultation.pdf)
  2. Based on results and experiences in the Hip and Knee Replacement Pilot study, a 15 day standard from referral to first consultation was deemed achievable by the Physician Steering Committee. Although during the Hip and Knee Replacement Pilot there were no surgery backlogs or day-to-day operational issues involved, implementation of the care path aims to ultimately eliminate those backlogs and a 15 day standard can be achieved. Timely assessment is important to patient care, thus it is important that we continue to monitor progress toward this goal. At this time however, based on literature investigating Maximum Acceptable Waiting Time (MAWT), 8-12 weeks was deemed acceptable. ('There are too many of us to fix.' Patients' views of acceptable waiting times for hip and knee replacement. Conner-Spady B., Sanmartin C., Johnston G., McGurran J., Kehler M., Noseworthy T. *EMBASE Journal of Health Services Research and Policy*. 14(4)(pp 212-218), 2009) *Clinical Committee Consensus, December 7, 2010*.
  3. A key principle of the care path states patients are required to participate in their care. Any surgery is stressful on the patient, and there are many things to remember and act upon. Additionally, each patient is required to designate a support person who will attend each clinic appointment and play an active role in preparing the patient for surgery, in achieving the planned length of stay, in transitioning the patient to home and in achieving desired recovery outcomes. A Surgical Patient Agreement has been developed to ensure patients and their support persons are informed of and, by signing, made accountable for arrangements contributing to optimal outcomes. Behavioural contracting is a strategy increasingly used by health professionals to improve patient compliance to health regimens. The literature provides evidence that contracts have been effective in promoting health behaviors by using reinforcement as a way to increase the likelihood that patients will follow instructions in order to reach agreed upon goals. *Clinical Committee Consensus, November 7, 2011*.
  4. Lemaigen, A., Armand-Lefevre, L., Birgand, G., Mabileau, G., Lolom, I., Ghodbane, W., ... Lucet, J.-C. (2018). Thirteen-year experience with universal *Staphylococcus aureus* nasal decolonization prior to cardiac surgery: a quasi-experimental study. *Journal of Hospital Infection*, 100(3), 322–328. doi: 10.1016/j.jhin.2018.04.023
  5. Hetem, D. J., Bootsma, M. C. J., & Bonten, M. J. M. (2015). Prevention of Surgical Site Infections: Decontamination With Mupirocin Based on Preoperative Screening for *Staphylococcus aureus* Carriers or Universal Decontamination? *Clinical Infectious Diseases*, 62(5), 631–636. doi: 10.1093/cid/civ990
  6. Safer Healthcare Now! Prevent Surgical Site Infections Getting Started Kit. 2014. <http://www.patientsafetyinstitute.ca/en/toolsResources/Documents/Interventions/Surgical%20Site%20Infection/SSI%20Getting%20Started%20Kit.pdf>
- According to a 2006 Cochrane Review titled “Preoperative hair removal to reduce surgical site infection”. The evidence finds no difference in surgical site infections (SSIs) among patients who have had hair removal prior to surgery and those who have not. If necessary to remove hair then both clipping and depilatory creams results in fewer SSIs than shaving with a razor.” Due to the incidence of reaction to the depilatory creams, hair removal as needed should be performed with a clipper prior to entering the surgical suite. *Clinical Committee Consensus, October 15, 2009*.

7. Alberta Bone and Joint Health Institute. *Prophylactics antibiotic use in hip and knee arthroplasty (evidence review)*. Available at: [ABJHI Evidence Review – Prophylactics Antibiotic Use](#)

Alberta Health Services. Bugs and Drugs: Antibiotics Prophylaxis Recommendation for Adult Joint Replacement. <http://www.bugsanddrugs.org/F90386B1-7292-4B28-AAB3-B51E69EE0626>

8. Alberta Health Services. Safe Surgery Checklist. Policy and Forms. 2014.

9. Fletcher N, Sofianos D, Berkes MB, Obremsky WT. Prevention of perioperative infection. *Journal of Bone and Joint Surgery - Series A* 2007 Jul;89(7):1605-1618.

Ostrander RV, Botte MJ, Brage ME. Efficacy of surgical preparation solutions in foot and ankle surgery. *Journal of Bone and Joint Surgery - Series A* 2005 May;87(5):980-985.

Saltzman MD, Nuber GW, Gryzlo SM, Marecek GS, Koh JL. Efficacy of surgical preparation solutions in shoulder surgery. *Journal of Bone and Joint Surgery - Series A* 2009 01 Aug;91(8):1949-1953.

10. A literature review completed in 2016 demonstrated that foley use increases the risk of urinary tract infections and reduces mobilization. Studies comparing patients who received catheters versus those who did not showed low in and out catheterization and no adverse events for those who did not. Therefore, foley catheters should not be routinely used in patients receiving elective arthroplasty and should only be used if appropriate for the patient (eg enlarged prostate). *Clinical Committee Consensus, October 27, 2016*.

Huang Z, Ma J, Shen B, Pei F. General Anesthesia: To catheterize or not? A prospective randomized controlled study of patients undergoing total knee surgery. *Journal of Arthroplasty* 30 (2015): 502-506.

Miller A, McKenzie J, Greenky M, Shaw E, Gandhi K, Hozack W, Parvizi J. Spinal Anesthesia: Should everyone receive a urinary catheter? A randomized, prospective study of patients undergoing total hip arthroplasty. *Journal of Bone and Joint Surgery* 95 (2013): 1498-503.

Tischler E, Restropo C, Oh J, Matthews C, Chen A, Parvizi J. Urinary retention is rare after total joint arthroplasty when using opioid-free regional anesthesia. *Journal of Arthroplasty* 31 (2016): 480-483.

11. Alberta Health Services. Surgical Site Verification and Marking. Policy and Forms. 2017.

Capital Health. Summary of Corporate Administrative Directive #2.6.6. (<http://www.intranet2.capitalhealth.ca/policies>) and UAH Patient Care Manual Policy #21.8 ([http://www.intranet2.capitalhealth.ca/uah-virtuallearningcentre/administrativepoliciesandprocedures/toc\\_care.htm](http://www.intranet2.capitalhealth.ca/uah-virtuallearningcentre/administrativepoliciesandprocedures/toc_care.htm))

12. Drains have not shown a clear advantage, represent an additional cost and expose patients to a high risk of transfusion (Walmsley et al., 2005). Based on Clinical Committee consensus and supporting evidence, the standard of care will be: "Hemovac drains – No Drains Hips or Knees" *Clinical Committee Consensus, October 15, 2009*

Walmsley, P. J., Kelly, M. B., Hill, R. M. F., & Brenkel, I. (2005). A prospective, randomised, controlled trial of the use of drains in total hip arthroplasty. *Journal of Bone and Joint Surgery - Series B*, 87(10), 1397-1401. .

13. Patient Blood Management Program AHS Calgary Zone. Protocol: Tranexamic Acid (TXA) in Arthroplasty Surgery. 2017.
14. Alberta Bone and Joint Health Institute. *Anticoagulation guidelines (rapid review)*.
15. Diabetes, Obesity and Nutrition Strategic Clinical Network. (2017). Inpatient Diabetes Management. <http://www.albertahealthservices.ca/scns/Page10970.aspx>
16. Ni, S. H., Jiang, W. T., Guo, L., Jin, Y. H., Jiang, T. L., Zhao, Y., Zhao, J. (2015). Cryotherapy on postoperative rehabilitation of joint arthroplasty. *Knee Surgery, Sports Traumatology Arthroscopy*. 23, 3354-3361.
17. Alberta Bone and Joint Health Institute. *New oral anticoagulants (evidence review)* . Available at: [ABJHI Evidence Review – New Oral Anticoagulants](#)
18. Alberta Bone and Joint Health Institute. *Perioperative use of NSAIDs by total hip and total knee arthroplasty patients (evidence review)*. Available at: [ABJHI Evidence Review - Perioperative use of NSAIDs](#)
19. This care path item has been updated to include the possible use of a Hip and Knee clinic discharge facilitator on site at acute care centers. It is imperative that variances from the surgical agreement are identified as early as possible by an individual who is fully aware of the patient's circumstances in terms of what is required to ensure optimal outcomes. The surgical patient agreement includes planned discharge date and location and planned functional and social requirements. *Clinical Committee Consensus, October 15, 2009*.
20. Canadian Orthopedic Association Consensus Statement: Patients with Total Joint Replacements having Dental Procedures. 2016. [http://www.coa-aco.org/images/stories/library/health\\_policy/Consensus\\_Statement\\_2016\\_Patients\\_with\\_Total\\_Joint\\_Replacements\\_having\\_Dental\\_Procedures.pdf](http://www.coa-aco.org/images/stories/library/health_policy/Consensus_Statement_2016_Patients_with_Total_Joint_Replacements_having_Dental_Procedures.pdf) (accessed October 2016).
21. The Clinical Committee has confirmed the importance of follow up appointments at two weeks, six weeks, twelve weeks and one year post surgery. At two weeks, a nurse at the clinic assesses progress, removes the staples, identifies the need for post-operative physiotherapy and enforces precautions. At six weeks, the surgeon assesses progress, orders x-rays if not previously completed and identifies any early concerns. At twelve weeks, the surgeon and care team re-assess progress, order x-rays if not previously completed and identify any concerns. Additionally, critical outcomes measures are to be collected at twelve weeks. At one year, surgical outcome is assessed and x-rays are reviewed.  
  
Monitoring is completed every five years at surgeon's discretion for primaries and annually at surgeon's discretion for revisions. Monitoring requires a review of current x-rays and an in-clinic assessment. If prosthesis problem/issue/failure is identified, follow-up should be expedited. *Clinical Committee Consensus, December 13, 2012*.
22. American Academy of Orthopaedic Surgeons. Diagnosis of Periprosthetic Joint Infections of the Hip and Knee.
23. Alberta Bone and Joint Health Institute. *Outcomes of primary and revision aseptic total hip and total knee arthroplasty with antibiotic-loaded bone cement (evidence review)*. Available at: [ABJHI Evidence Review – Outcomes of primary and revision aseptic total H&K with antibiotic-loaded bone cement](#)

24. Thomas W Wainwright, Mike Gill, David A McDonald, Robert G Middleton, Mike Reed, Opinder Sahota, Piers Yates & Olle Ljungqvist (2020) Consensus statement for perioperative care in total hip replacement and total knee replacement surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations, *Acta Orthopaedica*, 91:1, 3-19, DOI:10.1080/17453674.2019.1683790
25. Levy N, Quinlan J, El-Boghdadly K, Fawcett WJ, Agarwal V, Bastable RB, Cox FJ, de Boer HD, Dowdy SC, Hattingh K, Knaggs RD, Mariano ER, Pelosi P, Scott MJ, Lobo DN, Macintyre PE. An international multidisciplinary consensus statement on the prevention of opioid-related harm in adult surgical patients. *Anaesthesia*. 2021 Apr;76(4):520-536. doi: 10.1111/anae.15262. Epub 2020 Oct 7. PMID: 33027841
26. Hance A. Clarke, Varuna Manoo, Emily A. Pearsall, Akash Goel, Adina Feinberg, Aliza Weinrib, Jenny C. Chiu, Bansi Shah, Salima S. J. Ladak, Sarah Ward, Sanjho Srikandarajah, Savtaj S. Brar & Robin S. McLeod (2020) Consensus Statement for the Prescription of Pain Medication at Discharge after Elective Adult Surgery, *Canadian Journal of Pain*, 4:1, 67-85, DOI: [10.1080/24740527.2020.1724775](https://doi.org/10.1080/24740527.2020.1724775)
28. Anger, M., Valovska, T., Beloeil, H., Lirk, P., Joshi, G.P., Van de Velde, M., Raeder, J. and (2021), PROSPECT guideline for total hip arthroplasty: a systematic review and procedure-specific postoperative pain management recommendations. *Anaesthesia*, 76: 1082-1097. <https://doi.org/10.1111/anae.15498>
29. Registered Nurses' Association of Ontario: [Pasero Opioid-Induced Sedation Scale \(POSS\)](#)
30. [Michigan Opioid Prescribing Engagement Network \(OPEN\)](#)